arXiv:hep-th/9503081v1 12 Mar 1995

UPR-650-T

Addendum to Supersymmetric Dyonic Black Holes in Kaluza-Klein Theory

Mirjam Cvetič [*] and Donam Youm[†]
Physics Department
University of Pennsylvania, Philadelphia PA 19104-6396
(March 1995)

Abstract

We complete the study of 4-dimensional (4-d), static, spherically symmetric, supersymmetric black holes (BH's) in Abelian $(4+n)$-d Kaluza-Klein theory, by showing that for such solutions n electric charges $\vec{\mathcal{Q}} \equiv (Q_1, ..., Q_n)$ and n magnetic charges $\vec{\mathcal{P}} \equiv (P_1, ..., P_n)$ are subject to the constraint $\vec{\mathcal{P}} \cdot \vec{\mathcal{Q}} = 0$. All such solutions can be obtained by performing the $SO(n)$ rotations, which do not affect the 4-d space-time metric and the volume of the internal space, on the supersymmetric $U(1)_M \times U(1)_E$ BH's, *i.e.*, supersymmetric BH's with a diagonal internal metric.

In Ref. [1], 4-dimensional (4-d), static, spherically symmetric, supersymmetric black holes (BH's) in Abelian $(4+n)$-d Kaluza-Klein (KK) theory were obtained with a diagonal internal metric Ansatz. Such BH's correspond to $U(1)_M \times U(1)_E$ configurations, *i.e.*, they have at most one electric charge Q and one magnetic charge P which necessarily arise from different $U(1)$ groups. Here, we complete the study of 4-d static, spherically symmetric, supersymmetric BH's in Abelian KK theories by addressing the corresponding solutions with a general non-diagonal internal metric Ansatz. We show that n electric charges $\vec{\mathcal{Q}} \equiv (Q_1, ..., Q_n)$ and n magnetic charges $\vec{\mathcal{P}} \equiv (P_1, ..., P_n)$ of such BH's are subject to the constraint $\vec{\mathcal{P}} \cdot \vec{\mathcal{Q}} = 0$. All such solutions can be obtained by performing the global $SO(n)$ rotations [2] on the supersymmetric $U(1)_M \times U(1)_E$ BH solutions.

Throughout we use the notation specified in Ref. [1].

As discussed in Section 4.1 of Ref. [1], the supersymmetric BH configurations are solutions of the Killing spinor equations, which correspond to the vanishing of supersymmetric transformations on the dimensionally reduced $(4+n)$-d gravitino, *i.e.*, $\delta\psi^{\mathbf{m}}_{\mu} = \delta\psi^{\mathbf{m}}_{\tilde{\mu}} = 0$. Note, $\psi^{\mathbf{m}}_{\mu}$ and $\psi^{\mathbf{m}}_{\tilde{\mu}}$ ($\mathbf{m} = 1, \cdots, 2^{[\frac{n}{2}]}$, $\mu = 0, \cdots, 3$, $\tilde{\mu} = 4, \cdots, (n+3)$) are the corresponding 4-d gravitino(s) and dilatino(s), obtained by the dimensional reduction of $(4+n)$-d gravitino(s). (See Eqs. (3.6) and (3.7) of Ref. [1] for the explicit form of the corresponding supersymmetry transformations.)

---

[*] E-mail address: cvetic@cvetic.hep.upenn.edu

[†] E-mail address: youm@cvetic.hep.upenn.edu

For spherically symmetric configurations with the 4-d metric Ansatz given by Eq. (4.2) of Ref. [1] and, however, now with a general non-diagonal internal metric Ansatz, one can rewrite the t-component $\delta\psi^{\mathbf{m}}_{t} = 0$ of the 4-d gravitino Killing spinor equation, analogous to Eq. (4.9) in Ref. [1], as:

$$\frac{R}{\sqrt{\lambda}}\left(\partial_r\lambda - \frac{1}{\alpha}\lambda\partial_r\varphi\right)\gamma^{03}\varepsilon - \sum_{\tilde{a}=4}^{n+3}\tilde{\mathbf{Q}}^{\tilde{a}}(\gamma^{35}\otimes\gamma^{\tilde{a}})\varepsilon = 0 \ , \tag{1}$$

where $\tilde{\mathbf{Q}}^{\tilde{a}} \equiv \mathrm{e}^{-\frac{\alpha}{2}\varphi}(\Phi^{-1})^{\tilde{a}}_{\tilde{\pi}}Q^{\tilde{\pi}}$. Note, $\alpha = \sqrt{\frac{n+2}{n}}$, $\Phi^{\tilde{a}}_{\tilde{\pi}}$ ($\tilde{a}, \tilde{\pi} = 4, \cdots, n+3$) is the n-bein of the unimodular part of the internal metric, φ is the dilaton (the volume of the internal space), and λ and R are the (t,t) and (θ,θ) components of the 4-d metric $g_{\mu\nu}$, respectively. Here, $Q^{\tilde{\pi}}$ are electric charges obtained by integrating the Maxwell's equation $\partial_r(\mathrm{e}^{\alpha\varphi}R\lambda\Phi^{\tilde{a}}_{\tilde{\pi}}\Phi^{\tilde{a}}_{\tilde{\lambda}}F^{\tilde{\lambda}\ tr}) = 0$.

The θ-component $\delta\psi^{\mathbf{m}}_{\theta} = 0$ of the 4-d gravitino Killing spinor equation, analogous to Eq. (4.10), supplemented by the constraint Eq. (5.2) in Ref. [1], which is always satisfied by the Killing spinors of spherically symmetric configurations, assumes the following form:

$$\left[2\sqrt{R} - \sqrt{\lambda}\left(\partial_r R - \frac{1}{\alpha}R\partial_r\varphi\right)\right]\gamma^{13}\varepsilon - \sum_{\tilde{a}=4}^{n+3}\tilde{\mathbf{P}}^{\tilde{a}}(\gamma^{25}\otimes\gamma^{\tilde{a}})\varepsilon = 0 \ , \tag{2}$$

where $\tilde{\mathbf{P}}^{\tilde{a}} = \mathrm{e}^{\frac{\alpha}{2}\varphi}\Phi^{\tilde{a}}_{\tilde{\pi}}P^{\tilde{\pi}}$. Here, $P^{\tilde{\pi}}$ are magnetic charges satisfying the Maxwell's equation $\partial_\theta(\mathrm{e}^{\alpha\varphi}R\lambda\sin\theta\Phi^{\tilde{a}}_{\tilde{\pi}}\Phi^{\tilde{a}}_{\tilde{\lambda}}F^{\tilde{\lambda}\ \theta\phi}) = 0$ with $F^{\tilde{\lambda}}_{\theta\phi} = P^{\tilde{\lambda}}\sin\theta$.

Eqs. (1) and (2) can be satisfied if and only if the lower two-component spinors $\varepsilon^{\mathbf{m}}_{\ell}$ and the upper two-component spinors $\varepsilon^{\mathbf{m}}_{u}$ ($(\varepsilon^{\mathbf{m}})^T \equiv (\varepsilon^{\mathbf{m}}_{u}, \varepsilon^{\mathbf{m}}_{\ell})$) satisfy the following constraint:

$$\eta_Q\sum_{\tilde{a}=4}^{n+3}\tilde{\mathbf{Q}}'^{\ \tilde{a}}(\gamma^{\tilde{a}})^{\mathbf{m}}_{\ \mathbf{n}}\ \varepsilon^{\mathbf{n}}_{\ell} = \varepsilon^{\mathbf{m}}_{u} = i\eta_P\sum_{\tilde{a}=4}^{n+3}\tilde{\mathbf{P}}'^{\ \tilde{a}}(\gamma^{\tilde{a}})^{\mathbf{m}}_{\ \mathbf{n}}\ \varepsilon^{\mathbf{n}}_{\ell} \ , \quad \eta_{Q,P} = \pm 1 \ , \tag{3}$$

where $i \equiv \sqrt{-1}$. Here, $\tilde{\mathbf{Q}}'^{\ \tilde{a}} \equiv \tilde{\mathbf{Q}}^{\tilde{a}}/[R\lambda^{-1/2}(\partial_r\lambda - \frac{1}{\alpha}\lambda\partial_r\varphi)]$ and $\tilde{\mathbf{P}}'^{\ \tilde{a}} \equiv \tilde{\mathbf{P}}^{\tilde{a}}/[2\sqrt{R} - \sqrt{\lambda}(\partial_r R - \frac{1}{\alpha}R\lambda\partial_r\varphi)]$ satisfy the constraint:

$$\sum_{\tilde{a}=4}^{n+3}(\tilde{\mathbf{Q}}'^{\ \tilde{a}})^2 = \sum_{\tilde{a}=4}^{n+3}(\tilde{\mathbf{P}}'^{\ \tilde{a}})^2 = 1 \ . \tag{4}$$

Multiplying the left-most hand side of Eq. (3) by $\sum_{\tilde{b}=4}^{n+3}\tilde{\mathbf{P}}'^{\ \tilde{b}}(\gamma^{\tilde{b}})$ and the right-most hand side by $\sum_{\tilde{b}=4}^{n+3}\tilde{\mathbf{Q}}'^{\ \tilde{b}}(\gamma^{\tilde{b}})$, and summing the two resultant equations, along with the identity $\{\gamma^{\tilde{a}}, \gamma^{\tilde{b}}\} = -2\delta^{\tilde{a}\tilde{b}}$, one has the result:

$$\sum_{\tilde{a}=4}^{n+3}\tilde{\mathbf{P}}'^{\ \tilde{a}}\tilde{\mathbf{Q}}'^{\ \tilde{a}} = 0 \ , \tag{5}$$

or equivalently:

$$\sum_{\tilde{\pi}=4}^{n+3} P^{\tilde{\pi}} Q^{\tilde{\pi}} = \vec{\mathcal{P}} \cdot \vec{\mathcal{Q}} = 0 \ . \tag{6}$$

Therefore, supersymmetric BH's have constrained charge configurations with the electric charge vector $\vec{\mathcal{Q}}$ and the magnetic charge vector $\vec{\mathcal{P}}$ orthogonal to one another.

New supersymmetric BH solutions can be generated by performing the global $SO(n)$ rotations on known supersymmetric BH solutions. Namely, the effective 4-d KK Lagrangian density (see Eq. (3.4) of Ref. [1]) is invariant under the global $SO(n)$ transformations [2]:

$$\Phi^{\tilde{\lambda}}_{\tilde{a}} \to U^{\tilde{\lambda}}_{\tilde{\pi}} \Phi^{\tilde{\pi}}_{\tilde{a}} \ , \qquad A^{\tilde{\lambda}}_{\lambda} \to U^{\tilde{\lambda}}_{\tilde{\pi}} A^{\tilde{\pi}}_{\lambda} \ , \tag{7}$$

where U is an $n \times n$ matrix of the $SO(n)$ transformations. Transformations (7) affect the unimodular part of the internal metric and the gauge fields, however, they leave the 4-d space-time metric $g_{\mu\nu}$ and the dilaton φ intact. In addition, since the $SO(n)$ transformations are also the symmetry of the Killing spinor equations, the transformed solutions remain supersymmetric, *i.e.*, they also satisfy the Killing spinor equations.

The $SO(n)$ transformations (7) on the $U(1)_M \times U(1)_E$ supersymmetric BH's generate solutions with n electric $\vec{\mathcal{Q}}$ and n magnetic $\vec{\mathcal{P}}$ charges which are subject to the constraint $\vec{\mathcal{P}} \cdot \vec{\mathcal{Q}} = 0$ [2]. Therefore, those are all the 4-d static, spherically symmetric, supersymmetric BH's in Abelian KK theories.

ACKNOWLEDGMENTS

The work is supported by U.S. DOE Grant No. DOE-EY-76-02-3071, and the NATO collaborative research grant CGR 940870.

REFERENCES

[1] M. Cvetič and D. Youm, Nucl. Phys. **B438**, 182 (1995), hep-th # 9409119.
[2] M. Cvetič and D. Youm, UPR-645-T preprint (1994), hep-th # 9502099.